




RESIDUAL BROADLEAF AND GRASS CONTROL IN SUNFLOWERS


VELVETLEAF, LAMBSQUARTERS, MORNINGGLORIES, PIGWEEDS, KOCHIA

- Preemergence control of Kochia, Russian thistle, Pigweeds, Palmer pigweed, Lambsquarters, Waterhemp, and Morningglories.
- Also controls annual grasses: green, yellow, and giant foxtail, barnyard grass, and others.
- Greater crop flexibility with reduced rotation intervals on corn and other crops to allow fall application.
- Tank mix with any glyphosate for burndown program.
- Flexible, can be used in no-till or conventional tillage systems

Kochia Control in Sunflower - KSU



Redroot Pigweed Control in Sunflower - NDSU



Sunflower Use Information

Fall or Spring: Prior to planting up to 3 days after planting

Average Use Rates: 19-38 oz/A (See label for specific soil, organic matter use rate recommendations.)

- Coarse soils 19-25 oz/A
- Medium soils 19-32 oz/A
- Fine soils 21-38 oz/A



Rotational Crop Guideline

Soybean, Sunflower	Anytime
Wheat, Barley, Rye, Triticale	4 ½ months
Corn, Field & Seed	10 months
Sorghum	10 months
Alfalfa, Other cereal grains	12 months
Sweet Corn, Popcorn	18 months
Canola	12 months
Sugar Beets	36 months

Always read and follow directions. Spartan Elite Herbicide is not registered for sale or use in California. FMC and Spartan are trademarks of FMC Corporation or an affiliate. Dual II Magnum is a trademark of a Syngenta Group Company. Prowl is a trademark of BASF SE. Select is a registered trademark of Valent U.S.A. Corporation. ©2017 FMC Corporation. All rights reserved.